

Investigation of ^{10}Li resonance component in ^{11}Li via the $^{11}\text{Li}(p,d)$ reaction

Friday, 15 July 2016 11:45 (15 minutes)

The unbound system ^{10}Li is of great interest for the description of the structure of the Borromean neutron halo nucleus ^{11}Li [1,2]. Borromean neutron halo nuclei are unusual weakly bound states of a core nucleus plus two neutrons. No transfer reaction experiment has been done so far which directly looks at the sub-component ^{10}Li within ^{11}Li . While earlier measurements have indicated possible resonances in ^{10}Li [3-6], it is still not well established which resonance contributes to the ground state configuration of ^{11}Li and by what spectroscopic factor. To obtain such information a decisive way can be investigating the transfer of one-neutron from ^{11}Li .

The presentation will report observations on ^{10}Li studied through the first measurement of the $p(^{11}\text{Li},d)$ one-neutron transfer reaction at beam energy of 6A MeV. This was performed using a solid H2 target at the newly constructed IRIS facility at TRIUMF. The ^{10}Li residue was populated strongly as a resonance with energy $E_r = 0.62 \pm 0.04$ MeV having a total width $\Gamma = 0.33 \pm 0.07$ MeV. The angular distribution of this resonance is characterized by neutron occupying the $1p_{1/2}$ orbital. A DWBA analysis yields a spectroscopic factor of 0.67 ± 0.12 for $p_{1/2}$ removal strength from the ground state of ^{11}Li to the region of the peak.

The presentation will report observations on ^{10}Li studied through the first measurement of the $p(^{11}\text{Li},d)$ one-neutron transfer reaction at beam energy of 6A MeV. This was performed using a solid H2 target at the newly constructed IRIS facility at TRIUMF. The ^{10}Li residue was populated strongly as a resonance with energy $E_r = 0.62 \pm 0.04$ MeV having a total width $\Gamma = 0.33 \pm 0.07$ MeV. The angular distribution of this resonance is characterized by neutron occupying the $1p_{1/2}$ orbital. A DWBA analysis yields a spectroscopic factor of 0.67 ± 0.12 for $p_{1/2}$ removal strength from the ground state of ^{11}Li to the region of the peak.

- [1] E. Garrido, D.V. Fedorov, A.S. Jensen, Nucl. Phys. A 700 (2002) 117.
- [2] I. Tanihata, H. Savajols, R. Kanungo, Prog. in Part. and Nucl. Phys. 68 (2013) 215.
- [3] H.B. Jeppesen, A.M. Moro, U.C. Bergmann, et al., Phys. Lett. B 642 (2006) 449.
- [4] K.H. Wilcox, R.B. Weisenmiller, G.J. Wozniak, et al., Phys. Lett. B 59 (1975) 142.
- [5] A.I. Amelin, M.G. Gornov, Yu.B. Gurov, et al., Yad. Fiz. 52 (1990) 1231.
- [6] M. Zinser, F. Humbert, T. Nilsson, et al., Nuclear Phys. A 619 (1997) 151176.

Primary authors: Dr SANETULLAEV, Alisher (Saint Mary's University, TRIUMF, Inha University in Tashkent); Prof. KANUNGO, Rituparna (Saint Mary's University)

Co-authors: Mr GALLANT, Aaron (TRIUMF); Prof. CHEN, Alan (McMaster University); Prof. SHOTTER, Alan (University of Edinburgh); Dr ROJAS, Alexander (TRIUMF); Dr HADINIA, Baharak (University of Guelph); Dr DAVIDS, Barry (TRIUMF); Dr UNSWORTH, Carl (TRIUMF); Prof. ANDREOIU, Corina (Simon Fraser University); Dr MILLER, David (TRIUMF); Ms MCNEICE, E. (Saint Mary's University); Dr HACKMAN, Greg (TRIUMF); Prof. CHRISTIAN, Gregory (Texas AM University - Cyclotron Institute); Dr SAVAJOLS, Herve (GANIL); Dr THOMPSON, Ian (LLNL); Prof. TANIHATA, Isao (RCNP, Beihang University); Ms PURCELL, J. (Saint Mary's University); Mr RANDHAWA, Jaspreet (Saint Mary's University, Halifax); Dr FALLIS, Jennifer (TRIUMF); Dr LIGHTHALL, Jonathan (TRIUMF); Mr FORTIN, Julien (Saint Mary's University, University of Laval); Mr TANAKA, Junki (Research Center for Nuclear Physics); Dr ALCORTA, Martin (TRIUMF); Mr KEEFE, Matthew (Saint Mary's University); Dr GALINSKY, Naomi (TRIUMF); Prof. GARRETT, Paul (University of Guelph); Dr BENDER, Peter (TRIUMF); Dr VOSS, Philip (TRIUMF); Dr KRUECKEN, Reiner (TRIUMF); Dr ISHIMOTO, Shigeru (KEK); Dr ROGER, Thomas (GANIL); Dr WANG, Zhimin (TRIUMF)

Presenter: Dr SANETULLAEV, Alisher (Saint Mary's University, TRIUMF, Inha University in Tashkent)

Track Classification: New instrumentation for direct reaction studies of exotic nuclei