nEDM 2017



Contribution ID: 36

Type: Oral

Critical Spin Dressing

Thursday, 19 October 2017 10:40 (25 minutes)

It has long been proposed [1] that spin dressing [2] could be employed to realize a highly effective helium-3 nuclear precession co-magnetometer for a neutron electric dipole moment search. The proposal requires application of an intense, continuous, and far off-resonant oscillating magnetic field in such a way that the apparent Larmor precession frequency of both species is modified. Under appropriate conditions a desirable situation known as critical dressing is anticipated: the neutron and the helium-3 nucleus (or more generally, any two spin species) are expected to behave as if they had the same gyromagnetic ratio and hence should precess at the same rate in a static magnetic field.

Spin dressing has been studied in the context of the neutron [3], helium-3 [4], and a variety of other systems [5]. Critical dressing, however, has not previously been demonstrated. We will present results from recent NMR experiments in which simultaneous dressing of ¹H and ¹⁹F nuclei is studied, and in which critical dressing is observed. Insight gleaned from these experiments is expected to inform strategies for integrating critical dressing into a neutron electric dipole moment search.

[1] Phys Rep 237, 1 (1994)

[2] J Phys (France) 30, 153 (1969)

[3] Phys Rev Lett 58, 2047 (1987)

[4] e.g. Phys Rev A 85, 3 (2012); Phys Rev C, 76, 5: 051302, (2007); and Phys Rev C, 84, 2: 022501, (2011)

[5] e.g. Nature 471, 83 (2011) or Nature 476, 185 (2011)

Email

mhayden@sfu.ca

Funding Agency

NSERC

Primary author: Mr TAVAKOLI-DINANI, Reza (Department of Physics, Simon Fraser University)

Co-author: Prof. HAYDEN, Mike (Physics Department, Simon Fraser University)

Presenter: Mr TAVAKOLI-DINANI, Reza (Department of Physics, Simon Fraser University)

Session Classification: ThMo2

Track Classification: Experimental techniques (cryogenic, room temperature, crystal)