



Contribution ID: 1

Type: **not specified**

## Search for supersymmetry in final states with two same-sign or three leptons and jets using 13 TeV ATLAS data

*Saturday, 17 February 2018 11:15 (15 minutes)*

A search for strongly produced supersymmetric particles using signatures involving multiple energetic jets and either two isolated same-sign leptons ( $e$  or  $\mu$ ), or at least three isolated leptons, is presented. The analysis relies on the identification of  $b$ -jets and high missing transverse momentum to achieve good sensitivity. A data sample of proton–proton collisions at  $\sqrt{s} = 13$  TeV recorded with the ATLAS detector at the Large Hadron Collider in 2015 and 2016, corresponding to a total integrated luminosity of  $36.1 \text{ fb}^{-1}$ , is used for the search. No significant excess over the Standard Model prediction is observed. The results are interpreted in several simplified supersymmetric models featuring  $R$ -parity conservation or  $R$ -parity violation, extending the exclusion limits from previous searches. In models considering gluino pair production, gluino masses are excluded up to 1.87 TeV at 95% confidence level. When bottom squarks are pair-produced and decay to a chargino and a top quark, models with bottom squark masses below 700 GeV and light neutralinos are excluded at 95% confidence level.

**Primary author:** DUCU, Otilia (University of Montreal)

**Presenter:** DUCU, Otilia (University of Montreal)

**Session Classification:** Session #4