Contribution ID: 3

Higgs Boson Production via Quark mediated Gluon Fusion at High Energy at Next-to-Next-to-Leading Power in Quark Mass

Friday, 17 February 2023 11:45 (15 minutes)

In perturbative QCD, processes involving quark scattering provide the simplest way of studying non-Abelian scattering amplitudes. To that end, in this talk I will discuss our calculation for the Form Factor of the Higgs boson production via light quark mediated Gluon Fusion process in the high energy/small quark mass limit, where the leading contribution comes in the form of large double logarithms for each power of the strong coupling constant (α _s). We reproduced the leading coefficient (Next-to-Leading order in quark mass (m_q)) for the asymptotic expansion of the form factor in the high energy approximation; and extended the analysis to next coefficient which is at Next-to-Next-to-Leading order (O(m_q^3)). Three distinct sources of diagrams contribute to this coefficient at this order; the complete analytic result for which is presented at three loops. We also present all order analysis in the large-N_c limit of QCD as well as for the opposite Abelian limit. Our results provide a universal estimate of the light quark effects at the higher orders for both the Higgs boson production and decay processes.

Supervisor

Alexander Penin

Funding Agency

University of Alberta

Supervisor Email

penin@ualberta.ca

Your Email

smodi@ualberta.ca

Primary author: MODI, Sneh (University of Alberta)
Co-author: Prof. PENIN, Alexander
Presenter: MODI, Sneh (University of Alberta)
Session Classification: February 17 Morning Session

Track Classification: Electroweak and Higgs Physics