Contribution ID: 37

An Auxiliary ATLAS Detector to Enhance the Detection of Long Lived Particles

Thursday, 16 February 2023 19:45 (15 minutes)

Several models that provide solutions to the Standard Model's open problems allow for the possibility of long lived particles (LLPs) with lifetimes, in some cases, in excess of 10³ seconds. It is proposed that charged LLPs produced at the Large Hadron Collider could interact with the ATLAS detector, eventually coming to rest and decaying. Other LLPs, such as neutral particles, may also exit ATLAS and decay in flight just outside the detector. We propose a low cost scintillating detector to be placed above the ATLAS detector to measure LLP signals as they decay within ATLAS during technical stops and other "beam off" times, thus minimizing Standard Model background. The same detector would provide a decay volume outside of ATLAS to detect LLP decays in flight during normal operation. To illustrate the efficacy of the proposed detector we simulate the production, interaction, and decay of long lived particles within select models.

Supervisor

James Pinfold

Funding Agency

NSERC

Supervisor Email

jpinfold@ualberta.ca

Your Email

jmkelly1@ualberta.ca

Primary author: KELLY, Joseph Mitchell (University of Alberta)Presenter: KELLY, Joseph Mitchell (University of Alberta)Session Classification: February 16 Evening Session

Track Classification: Physics Beyond the Standard Model