

Sensitivity to the $X(3872)$ total width at the Belle II experiment

Thursday, 9 May 2019 17:45 (15 minutes)

The Belle II experiment at the SuperKEKB energy-asymmetric e^+e^- collider is a substantial upgrade of the B factory facility at the Japanese KEK laboratory. The design luminosity of the machine is $8 \times 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ and the Belle II experiment aims to record 50 ab^{-1} of data, a factor of 50 more than its predecessor. From February to July 2018, the machine has completed a commissioning run and main operation of SuperKEKB has started in March 2019. The $X(3872)$ is an exotic hadron candidate and studying the $X(3872)$ partial widths is a good probe for the internal structure of this hadronic state. However, in order to derive partial widths, a measurement of its total width is needed. The large Belle II data set will provide an ideal environment to measure the $X(3872)$ total width since it will be possible to use the $X(3872) \rightarrow D^0 \bar{D}^0 \pi^0$ decay, which has a better mass resolution than $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ used in earlier work. In this presentation, we will give an overview of the analysis and the expected sensitivity to the $X(3872)$ total width

Email

peruzzi@lnf.infn.it

Primary author: Prof. PERUZZI, Ida Marena (INFN-LNF)

Presenter: HIRATA, Hikari (Nagoya University)

Session Classification: Parallel session 1

Track Classification: Heavy non- $q\bar{q}$ Mesons and Pentaquarks