Contribution ID: 6

Angular analysis of $B_{(s)} \rightarrow K_1(\phi) ll$ decay modes

The LHCb experiment has reported discrepancy of $(2.2-2.4)\sigma$ in the μ/e ratio of $\bar{B} \to \bar{K}^* l^+ l^-$ process, R_{K^*} , which reinforce the hints of lepton nonuniversality observed in $B^+ \to K^+ l^+ l^-$ process. We investigate the analogous lepton non-universality ratios and other asymmetries in $B \to K_1 l^+ l^-$ and $B_s \to \phi l^+ l^-$ processes both in model dependent and independent approches. We constrain the new parameter space consistent with experimental limit on $Br(B_s \to ll)$, $Br(\bar{B} \to X_s ll)$, $Br(\bar{B}^0 \to \bar{K}^0 ll)$, $Br(\bar{B} \to K^{(*)} \nu_l \bar{\nu}_l)$, $Br(\bar{B} \to X_s \nu_l \bar{\nu}_l)$, R_K and R_{K^*} parameters. We then show the effects of new parameters on the branching ratios, forwardbackward asymmetries, CP violating parameters of $B_{(s)} \to K_1(\phi) l^+ l^-$ processes. As like $R_{K^{(*)}}$, we also check the existence of the violation of lepton universality in these decay modes. We observe that the analysis of B decays to axial vector mesons can also serve as a good tool to probe physics beyond the SM.

Email

suchismita8792@gmail.com

Primary author: Dr SAHOO, Suchismita (Physical Research Laboratory)

Presenter: Dr SAHOO, Suchismita (Physical Research Laboratory)

Track Classification: Rare Decays of Hadrons and Leptons